

## Comment on "Reverse Brazil Nut Problem: Competition between Percolation and Condensation"

In a recent paper [1] the Brazil nut problem (BNP), where particles with large diameters rise to the top when exposed to vertical shaking and the reverse Brasil nut problem (RBNP) where the large particles segregate to the bottom were investigated. In realistic situations these effects are driven or at least accompanied by global convection. Convection rolls are observed already in mono-disperse systems: apparently vibrations lead to temperature gradients with temperature defined through the average kinetic energy of the particles. In order to suppress convection the authors present Molecular Dynamics (MD) simulations for binary hard sphere mixtures at *fixed* temperature. It appears legitimate to first consider this simplified situation although it might be rather remote from reality. Depending on the mass and diameter ratios of the two hard sphere species, the simulations show pronounced BNP and RBNP behaviours. For an explanation a new *condensation versus percolation* driven segregation mechanism was put forward, which has attracted much attention in the press [2]. The purpose of this comment is to show that the proposed mechanism is incorrect. All the effects observed in the MD simulations can be understood by simple conventional thermodynamics.

Because in the solid phase particles may no longer exchange positions [3], we conjecture that demixing must happen already in the fluid phase. Therefore we consider a binary hard sphere mixture with species  $A$  and  $B$  having masses  $m_A > m_B$  and diameters  $d_A > d_B$  in the liquid phase with an appropriate equation of state (EOS), e.g. the Mansoori-Carnahan-Starling-Leland equation [5], with  $\xi_i = (n_A/n) d_A^i + (n_B/n) d_B^i$

$$P = nT \cdot Z(x_A, x_B), \quad Z(x_A, x_B) = \frac{1}{1-x} + \frac{3x}{(1-x)^2} \frac{\xi_1 \xi_2}{\xi_3} + \frac{x^2(3-x)}{(1-x)^3} \frac{\xi_2^3}{\xi_3^2}, \quad (1)$$

where  $n = n_A + n_B$  is the particle density and  $x = x_A + x_B = (\pi/6)d_A^3 n_A + (\pi/6)d_B^3 n_B$  the packing fraction. It is then straightforward to compute the Helmholtz free energy [4] under gravity

$$F = E - T \cdot S. \quad (2)$$

Variation of  $F$  with respect to the particle densities  $n_A$  and  $n_B$  for fixed particle numbers  $N_A$  and  $N_B$  leads to two coupled nonlinear equations which are solved numerically. In the limit  $N_A d_A^3 \ll N_B d_B^3$  and  $d_A \gg d_B$  these equations respect Archimedes' principle. Note however, that the initial condition of equal layer numbers  $N_A d_A^2 = N_B d_B^2$  used in [1] is incompatible with this limit. Consequently Archimedes' principle need not be obeyed in the presented MD simulations in contrast to our naïve expectation. Calculated filling fractions  $x_A(z)$  and  $x_B(z)$  as functions of the height  $z$  are depicted in Fig. 1 for various mass ratios  $m_A/m_B$  keeping the diameter ratio  $d_A/d_B = 2$  and particle number ratio  $N_A/N_B = 1/4$  fixed. The density profiles compare very well with the MD simulations presented in [1]. For  $m_A/m_B = 2$  we encounter the BNP with almost pure B-liquid at the bottom. The  $m_A/m_B = 6$  case corresponds to the RBNP with an incomplete demixing at the bottom. For  $m_A/m_B = 4$  the two liquids remain mixed. All these details agree with the corresponding MD simulations. But note that in our calculation the temperatures  $T > T_C \geq T_A$  lie above the temperature  $T_C$  where the binary liquid begins to solidify: the total packing fraction lies sufficiently below close packing everywhere. In contrast the temperatures used in [1] are supposed to be quenched between  $T_B < T < T_A$  ( $T_A$  and  $T_B$  represent the temperatures where condensation sets in for the separated species [3], see below) such that parts of the sample are "frozen".




Figure 1: Calculated filling fractions  $x_A$  and  $x_B$  of binary hard sphere liquids under gravity for various mass ratios. The length unit equals the container size used in the MD simulations of Ref. [1]. The BNP and RBNP are clearly recognized. With decreasing temperature the profiles are shifted to smaller heights  $z$ .

We must conclude that demixing indeed occurs in the liquid phase. It is clearly due to the competition between *gravity and entropy* rather than to that of *condensation and percolation*. There is nothing mysterious or special going on and all the effects are understood by conventional thermodynamics. Finally, if the temperature is further decreased, the binary liquid eventually begins to condensate locally according to its filling fractions  $x_A$  and  $x_B$  in the fluid phase. In the RBNP this starts from the bottom whereas in the BNP, depending on the parameters, species  $A$  in the middle may condensate first (cf. Fig. 1). However, there exists no condensation driven demixing.

In order to discuss the transition from the BNP to the RBNP and also to understand the effects on an even simpler basis, we may consider the two species separated:  $A$  above  $B$  for the BNP and  $B$  above  $A$  for the RBNP. The advantage is that we may then use the much simpler EOS for one-component hard sphere fluids e.g. the Carnahan-Starling expression [6]

$$P = n T Z(x), \quad Z(x) = \frac{1 + x + x^2 - x^3}{(1 - x)^3} \quad (3)$$

obtained from (1) by setting  $d_A = d_B$ . The details of the EOS are unimportant here, in fact, every  $Z(x)$  which grows sufficiently fast with  $x$  (eigen-volume effect) leads to similar results. We assume that at a filling fraction  $x_0 \simeq 0.637$  equal to *random* close packing [7], instead of FCC or HCP close packing  $x_0 = \pi/\sqrt{18} \simeq 0.74048$ , condensation sets in, which leads to temperatures

$$T_A = \frac{\pi}{6} \frac{N_A m_A d_A^3}{x_0 Z(x_0)} \frac{g}{\text{area}}, \quad T_B = \frac{\pi}{6} \frac{N_B m_B d_B^3}{x_0 Z(x_0)} \frac{g}{\text{area}} \quad (4)$$

(gravity  $g$ ) which exceed those in [3] by roughly a factor of 3. We choose  $x_A(0) = x_0$  at the bottom of the RBNP which corresponds to the temperature  $T_{RBNP} = T_A + (d_A/d_B)^3 T_B > T_A + T_B$  where this system is just still fluid. Then also the BNP configuration is fluid, because there "freezing" sets in at lower temperatures (either at  $T_{BNP} = T_A$  at the boundary of the two species or at  $T_{BNP} = (d_B/d_A)^3 T_A + T_B$  at the bottom, depending on the parameters used). With the above choice the filling fractions at the bottom and at the boundaries are fixed by the corresponding pressures and the Helmholtz free energies  $F_{BNP}$  and  $F_{RBNP}$  may be computed. For  $F_{BNP} < F_{RBNP}$  the free energy favors the BNP and vice versa. We consider the curves  $F_{BNP} = F_{RBNP}$  which separate BNP from RBNP for the following cases:

- (i) equal number of particles  $N_A = N_B$
- (ii) equal number of layers  $N_A d_A^2 = N_B d_B^2$
- (iii) equal volumes  $N_A d_A^3 = N_B d_B^3$
- (iv) equal masses  $N_A m_A = N_B m_B$ .

The last case is not explicitly shown in Fig.2 because it almost coincides with case (ii). The MD simulation results, case (ii), are also shown with circles (BNP), boxes (RBNP) and triangles (mixed). In the vicinity of the curves mixing is expected, demixing occurs for parameter combinations lying sufficiently far away. We find that the particle numbers are of minor influence. This is expected: the BNP and RBNP should not be easily reversed by adding or subtracting a few particles of either species.




Figure 2: Curves which separate BNP from RBNP for different particle number ratios in the mass ratio and diameter ratio parameter space. The MD simulation results of Ref. [1] are also shown.

Lets contrast our results with the "percolation-condensation" formula

$$\frac{T_A}{T_B} = \frac{N_A m_A d_A^3}{N_B m_B d_B^3} = \frac{d_A^3}{d_B^3} \quad (5)$$

of Ref. [1]. This formula seems to give reasonable results  $d_A/d_B = (m_A/m_B)^{1/2}$  for the equal-number-of-layers case (ii), but certainly a straight line would not be worse. For equal volumes, case (iii), (5) leads to  $d_A/d_B = (m_A/m_B)^{1/3}$  and the  $m_A/m_B = 6$ ,  $d_A/d_B = 2$  system is predicted BNP, which is incorrect. The calculated profiles look very similar to the corresponding ones of case (ii) (Fig. 1) with the only difference that  $x_B$  is somewhat larger because of the relatively larger particle number  $N_B$ . An MD calculation certainly would confirm this. For equal particle numbers, case (i), the BNP is obtained for  $m_A < m_B$  whereas the equal mass case (vi) is predicted to be always mixed. All these implications are wrong and we must conclude that the formula (5) works only approximately for case (ii) by pure accident.

Preparing this comment, I realized that J.A. Both together with one of the authors of Ref.[1], D.C. Hong, have just submitted a paper to the archive [8], which treats the problem along similar lines. This paper, however, lacks the clear statement that the "condensation - percolation theory" is incorrect. In view of the publicity given to this picture [2] such a statement would seem appropriate.

H. Walliser

*Fachbereich Physik, Universität Siegen, D-57068 Siegen, Germany*

## References

- [1] D.C. Hong, P.V. Quinn and S. Luding, Phys. Rev. Lett. **86**, 3423 (2001).
- [2] "Shaking up is hard to do" by P. Ball, Nature Science Update, 23 march 2001; "Cracking the Brazil nut problem", Physics Web News, 11 april 2001; "This theory is nuts" by S. Reucroft and J. Swain, 17 april 2001, The Boston Globe; "Das Paranus-Problem" by M. Rauner in Die Zeit 33/2001.
- [3] D.C. Hong, Physica A **271**, 192 (1999); J.A.Both and D.C. Hong, Phys. Rev. E **64**, 061105 (2001).
- [4] S.B. Yuste, A. Santos and M. López de Haro, Europhys. Lett. **52**, 158 (2000).
- [5] G.A. Mansoori, N.F. Carnahan, K.E. Starling and T.W. Leland, J. Chem. Phys. **54**, 1523 (1971).
- [6] N.F. Carnahan and K.E. Starling, J. Chem. Phys. **51**, 635 (1961).
- [7] S. Torquato, T.M. Truskett and P.G. Debenedetti, Phys. Rev. Lett. **84**, 2064 (2000).
- [8] J.A. Both and D.C. Hong, cond-mat/0111347.